YouGov - Prostate cancer

Sample Size: 1456 adults in GB who were described as male at birth (i.e. who have a prostate)

| | | Ger | nder | Age | | | Social | Grade | Region | | | | | | | |
|---|-------|------|--------|-------|-------|-------|--------|-------|--------|------|------|--------|------------------|---------------------|-------|----------|
| | Total | Male | Female | 18-29 | 30-39 | 40-49 | 50-59 | 60-69 | 70-79 | ABC1 | C2DE | London | Rest of South | Midlands / Wales | North | Scotland |
| Weighted Sample | 1618 | 1606 | 12 | 226 | 253 | 385 | 273 | 237 | 219 | 899 | 720 | 195 | 492 | 370 | 405 | 157 |
| Unweighted Sample | 1456 | 1444 | 12 | 119 | 207 | 321 | 281 | 252 | 247 | 850 | 606 | 145 | 450 | 351 | 372 | 138 |
| | % | % | % | % | % | % | % | % | % | % | % | % | % | % | % | % |
| What symptoms of prostate cancer are you aware of? | | | | | | | | | | | | | | | | |
| Please list as many as you know. | | | | | | | | | | | | | | | | |
| If you do not know any, please select the 'don't | | | | | | | | | | | | | | | | |
| know' option. | | | | | | | | | | | | | | | | |
| Having to/feeling like you have to urinate more frequently | 13 | 13 | 0 | 5 | 6 | 11 | 20 | 18 | 19 | 15 | 10 | 14 | 12 | 16 | 11 | 15 |
| Difficulty urinating | 10 | 10 | 7 | 6 | 6 | 8 | 10 | 13 | 19 | 12 | 8 | 12 | 10 | 10 | 10 | 7 |
| Blood in urine | 6 | 6 | 0 | 5 | 3 | 6 | 6 | 11 | 9 | 7 | 5 | 6 | 5 | 9 | 6 | 6 |
| Other | 6 | 6 | 22 | 6 | 8 | 5 | 6 | 6 | 5 | 7 | 4 | 6 | 7 | 6 | 5 | 6 |
| Blood in stool | 3 | 3 | 10 | 2 | 3 | 4 | 3 | 4 | 1 | 3 | 3 | 1 | 3 | 3 | 3 | 7 |
| Low urine flow / poor stream | 3 | 3 | 0 | 0 | 1 | 2 | 4 | 7 | 4 | 3 | 2 | 1 | 3 | 3 | 3 | 2 |
| Pain (unspecified) | 3 | 3 | 7 | 3 | 3 | 4 | 2 | 2 | 2 | 3 | 3 | 2 | 2 | 2 | 4 | 4 |
| Painful urination | 2 | 2 | 0 | 2 | 1 | 2 | 1 | 2 | 3 | 3 | 1 | 5 | 1 | 1 | 2 | 2 |
| Weight loss | 1 | 1 | 10 | 2 | 1 | 2 | 1 | 1 | 0 | 2 | 1 | 2 | 2 | 0 | 1 | 2 |
| Specifically mention 'at night' re: more frequent urination | 1 | 1 | 0 | 0 | 0 | 0 | 2 | 4 | 2 | 2 | 1 | 0 | 1 | 2 | 1 | 3 |
| Enlarged prostate | 1 | 1 | 0 | 1 | 0 | 1 | 2 | 2 | 2 | 2 | 0 | 1 | 2 | 1 | 1 | 2 |
| Blood/Bleeding (unspecified) | 1 | 1 | 13 | 0 | 0 | 2 | 2 | 1 | 1 | 1 | 2 | 2 | 2 | 1 | 1 | 0 |
| Incontinence | 1 | 1 | 0 | 1 | 0 | 0 | 1 | 3 | 3 | 2 | 1 | 2 | 1 | 0 | 2 | 0 |
| Can't empty bladder | 1 | 1 | 0 | 0 | 0 | 1 | 2 | 1 | 1 | 1 | 1 | 3 | 1 | 0 | 1 | 1 |
| High PSA levels | 1 | 1 | 0 | 0 | 0 | 0 | 1 | 2 | 3 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| Back pain | 1 | 1 | 0 | 0 | 0 | 1 | 0 | 1 | 1 | 1 | 1 | 1 | 0 | 1 | 1 | 0 |
| Loss of appetite | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 1 | 1 |
| Blood in semen | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 |
| Bone pain | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Needing to rush to the toilet | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 |
| Straining or taking a long time while peeing | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Testicular pain | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Don't know | 68 | 68 | 58 | 79 | 78 | 70 | 62 | 60 | 54 | 63 | 74 | 67 | 68 | 65 | 70 | 68 |
| N/A | 1 | 1 | 0 | 0 | 0 | 0 | 1 | 0 | 3 | 1 | 1 | 0 | 1 | 1 | 0 | 0 |

YouGov - Prostate cancer

Sample Size: 1456 adults in GB who were described as male at birth (i.e. who have a prostate)

| | | As far as you are aware, what is the typical test prostate cancer? Please select all that apply. | | | | | |
|--|---------|--|---------------|------------------|--|--|--|
| | Total | A blood analysis test | A rectal exam | Both are typical | | | |
| Weighted Sample | 1618 | 144 | 535 | 517 | | | |
| Unweighted Sample | 1456 | 147 | 450 | 499 | | | |
| | % | % | % | % | | | |
| What symptoms of prostate cancer are you aware of? | | | | | | | |
| Please list as many as you know. | | | | | | | |
| If you do not know any, please select the 'don't | | | | | | | |
| know' option. | | | | | | | |
| Having to/feeling like you have to urinate more frequently | 13 | 17 | 13 | 20 | | | |
| Difficulty urinating | 10 | 15 | 8 | 16 | | | |
| Blood in urine | 6 | 11 | 5 | 10 | | | |
| Other | 6 | 5 | 5 | 9 | | | |
| Blood in stool | 3 | 2 | 4 | 4 | | | |
| Low urine flow / poor stream | 3 | 6 | 3 | 3 | | | |
| Pain (unspecified) | 3 | 2 | 4 | 3 | | | |
| Painful urination | 2 | 2 | 2 | 3 | | | |
| Weight loss | 1 | 1 | 1 | 3 | | | |
| Specifically mention 'at night' re: more frequent urination | 1 | 2 | 2 | 2 | | | |
| Enlarged prostate | 1 | 1 | 1 | 3 | | | |
| Blood/Bleeding (unspecified) | 1 | 2 | 2 | 1 | | | |
| Incontinence | 1 | 1 | 1 | 3 | | | |
| Can't empty bladder | 1 | 0 | 2 | 1 | | | |
| High PSA levels | 1 | 4 | 0 | 2 | | | |
| Back pain | 1 | 0 | 1 | 2 | | | |
| Loss of appetite | 0 | 1 | 1 | 0 | | | |
| Blood in semen | 0 | 0 | 0 | 1 | | | |
| Bone pain | 0 | 0 | 0 | 0 | | | |
| Needing to rush to the toilet | 0 | 1 | 0 | 0 | | | |
| Straining or taking a long time while peeing | 0 | 0 | 0 0 | 0 | | | |
| Testicular pain Don't know | 0 68 | 0 | 0 68 | 0 54 | | | |
| Don't know N/A | 68 1 | 53 2 | 68 0 | 54 1 | | | |
| N/A | T | | U | Т | | | |

YouGov - Prostate cancer

Sample Size: 1456 adults in GB who were described as male at birth (i.e. who have a prostate)

| | | Ger | nder | Age | | | Social | Grade | Region | | | | | | | |
|--|----------|------|----------|-------|----------|-------|--------|----------|----------|----------|------|----------|------------------|---------------------|----------|----------|
| | Total | Male | Female | 18-29 | 30-39 | 40-49 | 50-59 | 60-69 | 70-79 | ABC1 | C2DE | London | Rest of South | Midlands / Wales | North | Scotland |
| Weighted Sample | 1618 | 1606 | 12 | 226 | 253 | 385 | 273 | 237 | 219 | 899 | 720 | 195 | 492 | 370 | 405 | 157 |
| Unweighted Sample | 1456 | 1444 | 12 | 119 | 207 | 321 | 281 | 252 | 247 | 850 | 606 | 145 | 450 | 351 | 372 | 138 |
| | % | % | % | % | % | % | % | % | % | % | % | % | % | % | % | % |
| Have you ever had a test for prostate cancer? | | | | | | | | | | | | | | | | |
| Yes I have | 20 | 20 | 17 | 3 | 1 | 8 | 28 | 35 | 52 | 21 | 19 | 13 | 24 | 20 | 20 | 18 |
| No I have not | 80 | 80 | 83 | 97 | 99 | 92 | 72 | 65 | 48 | 79 | 81 | 87 | 76 | 80 | 80 | 82 |
| How long ago was your most recent prostate cancer test? | | | | | | | | | | | | | | | | |
| [Asked to those who have had a test for prostate cancer; n=352] | | | | | | | | | | | | | | | | |
| In the last 6 months | 19 | 18 | 59 | 0 | 0 | 23 | 18 | 24 | 16 | 19 | 18 | 19 | 20 | 19 | 16 | 19 |
| In the last year | 9 | 9 | 0 | 0 | 0 | 9 | 6 | 13 | 9 | 8 | 12 | 13 | 8 | 6 | 12 | 14 |
| In the last 18 months | 7 | 7 | 41 | 32 | 0 | 10 | 7 | 9 | 4 | 8 | 6 | 7 | 9 | 9 | 6 | 3 |
| In the last 2 years | 17 | 17 | 0 | 68 | 0 | 17 | 17 | 12 | 19 | 15 | 19 | 20 | 19 | 18 | 16 | 3 |
| In the last 2 and a half years | 6 | 6 | 0 | 0 | 0 | 6 | 10 | 4 | 5 | 4 | 8 | 0 | 6 | 6 | 8 | 4 |
| In the last 3 years | 6 | 6 | 0 | 0 | 0 | 4 | 5 | 6 | 8 | 8 | 3 | 8 | 6 | 8 | 3 | 10 |
| In the last 5 years | 16 | 16 | 0 | 0 | 0 | 15 | 18 | 17 | 16 | 18 | 14 | 9 | 16 | 12 | 21 | 22 |
| In the last 10 years | 11 | 11 | 0 | 0 | 55 | 4 | 9 | 9 | 14 | 14 | 8 | 7 | 10 | 16 | 10 | 12 |
| Don't know/can't recall | 8 | 8 | 0 | 0 | 45 | 11 | 9 | 6 | 8 | 6 | 11 | 17 | 6 | 6 | 8 | 13 |
| How willing would you be to have a routine test for prostate cancer? [Asked to those who have never had a test for prostate cancer; n=1104] | | | | | | | | | | | | | | | | |
| Very willing | 33 | 33 | 50 | 19 | 26 | 34 | 37 | 43 | 51 | 35 | 30 | 30 | 33 | 38 | 29 | 34 |
| Fairly willing | 39 | 39 | 30 30 | 44 | 20 47 | 41 | 31 | 43 32 | 30 | 40 | 38 | 42 | 33 40 | 35 | 40 | 34 39 |
| TOTAL WILLING | 39 72 | 72 | 30 80 | 62 | 73 | 75 | 67 | 52 75 | 30 81 | 40 75 | 68 | 42 72 | 73 | 73 | 40 69 | 73 |
| Not very willing | 12 | 12 | 0 | 15 | 11 | 9 | 16 | 10 | 8 | 13 | 10 | 15 | 12 | 10 | 12 | 8 |
| Not at all willing | 6 | 6 | 20 | 10 | 4 | 6 | 4 | 7 | 10 | 3 | 10 | 2 | 5 | 7 | 8 | 9 |
| TOTAL NOT WILLING | 18 | 18 | 20 | 24 | 15 | 15 | 20 | 17 | 18 | 16 | 20 | 17 | 17 | 16 | 20 | 17 |
| Don't know | 10 | 10 | 0 | 13 | 12 | 10 | 13 | 8 | 1 | 9 | 12 | 11 | 10 | 11 | 11 | 9 |
| Dontailou | | | - | | | | | ÷ | | Ŭ | | | | •• | | ~ |

YouGov - Prostate cancer

Sample Size: 1456 adults in GB who were described as male at birth (i.e. who have a prostate)

| | | As far as you are prostate cane | | |
|--|-------|---------------------------------|---------------|------------------|
| | Total | A blood analysis test | A rectal exam | Both are typical |
| Weighted Sample | 1618 | 144 | 535 | 517 |
| Unweighted Sample | 1456 | 147 | 450 | 499 |
| | % | % | % | % |
| Have you ever had a test for prostate cancer? | | | | |
| Yes I have | 20 | 38 | 17 | 31 |
| No I have not | 80 | 62 | 83 | 69 |
| How long ago was your most recent prostate cancer test? | | | | |
| [Asked to those who have had a test for prostate cancer; n=352] | | | | |
| In the last 6 months | 19 | 26 | 3 | 23 |
| In the last year | 9 | 8 | 8 | 11 |
| In the last 18 months | 7 | 8 | 9 | 7 |
| In the last 2 years | 17 | 25 | 15 | 16 |
| In the last 2 and a half years | 6 | 6 | 6 | 5 |
| In the last 3 years | 6 | 7 | 7 | 5 |
| In the last 5 years | 16 | 4 | 26 | 16 |
| In the last 10 years | 11 | 8 | 13 | 12 |
| Don't know/can't recall | 8 | 6 | 13 | 5 |
| How willing would you be to have a routine test for prostate cancer? [Asked to those who have never had a test for prostate cancer; n=1104] | | | | |
| Very willing | 33 | 48 | 31 | 43 |
| Fairly willing | 39 | 39 | 42 | 38 |
| TOTAL WILLING | 72 | 87 | 73 | 81 |
| Not very willing | 12 | 4 | 17 | 10 |
| Not at all willing | 6 | 5 | 6 | 3 |
| TOTAL NOT WILLING | 18 | 10 | 22 | 13 |
| Don't know | 10 | 3 | 5 | 6 |

YouGov - Prostate cancer

Sample Size: 1456 adults in GB who were described as male at birth (i.e. who have a prostate)

| | | Ger | nder | Age | | | Social | Grade | Region | | | | | | | |
|---|---------------|---------------|----------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|------------------|---------------------|---------------|---------------|
| | Total | Male | Female | 18-29 | 30-39 | 40-49 | 50-59 | 60-69 | 70-79 | ABC1 | C2DE | London | Rest of South | Midlands / Wales | North | Scotland |
| Weighted Sample | 1618 | 1606 | 12 | 226 | 253 | 385 | 273 | 237 | 219 | 899 | 720 | 195 | 492 | 370 | 405 | 157 |
| Unweighted Sample | 1456 | 1444 | 12 | 119 | 207 | 321 | 281 | 252 | 247 | 850 | 606 | 145 | 450 | 351 | 372 | 138 |
| | % | % | % | % | % | % | % | % | % | % | % | % | % | % | % | % |
| As far as you are aware, what is the typical test for prostate cancer? | | | | | | | | | | | | | | | | |
| A blood analysis test | 9 | 9 | 15 | 4 | 5 | 8 | 11 | 14 | 13 | 9 | 8 | 9 | 9 | 10 | 9 | 6 |
| A rectal exam (i.e. a doctor inserting their finger into the rectum to feel the prostate) | 33 | 33 | 23 | 48 | 40 | 33 | 28 | 28 | 22 | 33 | 33 | 28 | 31 | 36 | 35 | 35 |
| Both are typical | 32 | 32 | 36 | 17 | 24 | 31 | 36 | 40 | 44 | 34 | 29 | 33 | 35 | 28 | 32 | 31 |
| Neither are typical/Some other test is typical | 2 | 2 | 10 | 5 | 1 | 2 | 1 | 1 | 4 | 2 | 2 | 0 | 2 | 2 | 3 | 4 |
| Don't know | 24 | 24 | 16 | 26 | 30 | 26 | 24 | 16 | 17 | 21 | 28 | 30 | 24 | 23 | 22 | 24 |
| For each of the following, please say whether you think they are true or false. If you don't know, please choose the 'don't know' option. Prostate cancer is more common among younger males than older males TRUE FALSE Don't know | 4 64 32 | 4 64 32 | 23 55 22 | 8 56 36 | 6 60 34 | 4 65 31 | 3 67 30 | 3 70 27 | 2 68 30 | 3 70 26 | 5 56 38 | 3 68 29 | 4 63 33 | 6 62 32 | 3 66 31 | 4 65 31 |
| A healthy diet can reduce your risk of getting prostate cancer | | | | | | | | | | | | | | | | |
| TRUE | 59 | 59 | 48 | 63 | 65 | 63 | 61 | 57 | 45 | 64 | 54 | 59 | 58 | 63 | 59 | 58 |
| FALSE | 8 | 8 | 10 | 3 | 5 | 6 | 8 | 9 | 15 | 7 | 8 | 6 | 7 | 11 | 8 | 3 |
| Don't know | 33 | 33 | 42 | 33 | 30 | 31 | 31 | 34 | 40 | 29 | 38 | 35 | 35 | 27 | 33 | 39 |
| The NHS asks males to come in for prostate cancer tests once they reach a certain age, like it does for females and breast cancer | | | | | | | | | | | | | | | | |
| TRUE | 36 | 36 | 57 | 60 | 59 | 42 | 25 | 18 | 10 | 37 | 36 | 41 | 31 | 39 | 36 | 41 |
| FALSE | 30 | 30 | 34 | 10 | 8 | 19 | 32 | 54 | 62 | 33 | 25 | 24 | 32 | 28 | 30 | 35 |
| Don't know | 34 | 34 | 9 | 30 | 34 | 39 | 43 | 28 | 28 | 30 | 39 | 35 | 37 | 33 | 34 | 24 |

YouGov - Prostate cancer

Sample Size: 1456 adults in GB who were described as male at birth (i.e. who have a prostate)

| | | | e aware, what is the typical test for er? Please select all that apply. | | | | |
|---|-------|-----------------------|--|------------------|--|--|--|
| | Total | A blood analysis test | A rectal exam | Both are typical | | | |
| Weighted Sample | 1618 | 144 | 535 | 517 | | | |
| Unweighted Sample | 1456 | 147 | 450 | 499 | | | |
| | % | % | % | % | | | |
| As far as you are aware, what is the typical test for prostate cancer? | | | | | | | |
| A blood analysis test | 9 | 100 | 0 | 0 | | | |
| A rectal exam (i.e. a doctor inserting their finger into the rectum to feel the prostate) | 33 | 0 | 100 | 0 | | | |
| Both are typical | 32 | 0 | 0 | 100 | | | |
| Neither are typical/Some other test is typical | 2 | 0 | 0 | 0 | | | |
| Don't know | 24 | 0 | 0 | 0 | | | |
| you think they are true or false. If you don't know, please choose the 'don't know' option. Prostate cancer is more common among younger males than older males TRUE | 4 | 8 | 4 | 4 | | | |
| FALSE | 64 | 71 | 70 | 77 | | | |
| Don't know | 32 | 22 | 26 | 19 | | | |
| A healthy diet can reduce your risk of getting prostate cancer | | | | | | | |
| TRUE | 59 | 62 | 63 | 67 | | | |
| FALSE | 8 | 12 | 7 | 10 | | | |
| Don't know | 33 | 26 | 31 | 23 | | | |
| The NHS asks males to come in for prostate cancer tests once they reach a certain age, like it does for females and breast cancer | | | | | | | |
| TRUE | 36 | 27 | 51 | 36 | | | |
| FALSE | 30 | 43 | 24 | 37 | | | |
| Don't know | 34 | 29 | 25 | 26 | | | |
| | | • | | | | | |

YouGov - Prostate cancer

Sample Size: 1456 adults in GB who were described as male at birth (i.e. who have a prostate)

| | | Ger | ıder | Age | | | Social | Grade | Region | | | | | | | |
|---|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|------------------|---------------------|---------|----------|
| | Total | Male | Female | 18-29 | 30-39 | 40-49 | 50-59 | 60-69 | 70-79 | ABC1 | C2DE | London | Rest of South | Midlands / Wales | North | Scotland |
| Weighted Sample | 1618 | 1606 | 12 | 226 | 253 | 385 | 273 | 237 | 219 | 899 | 720 | 195 | 492 | 370 | 405 | 157 |
| Unweighted Sample | | 1444 | 12 | 119 | 207 | 321 | 281 | 252 | 247 | 850 | 606 | 145 | 450 | 351 | 372 | 138 |
| | % | % | % | % | % | % | % | % | % | % | % | % | % | % | % | % |
| Vulnerability to prostate cancer is genetic i.e. you are more like to get prostate cancer if other members of your family got prostate cancer | | | | | | | | | | | | | | | | |
| TRUE | 35 | 35 | 17 | 41 | 35 | 38 | 33 | 29 | 32 | 37 | 31 | 43 | 36 | 34 | 31 | 31 |
| FALSE | 14 | 14 | 11 | 8 | 11 | 11 | 12 | 25 | 16 | 14 | 14 | 11 | 12 | 17 | 16 | 12 |
| Don't know | 52 | 51 | 72 | 51 | 54 | 51 | 56 | 46 | 52 | 49 | 54 | 46 | 53 | 49 | 53 | 57 |
| Prostate cancer requires immediate treatment once discovered | | | | | | | | | | | | | | | | |
| TRUE | 66 | 66 | 71 | 55 | 65 | 70 | 66 | 65 | 70 | 65 | 67 | 62 | 64 | 71 | 65 | 67 |
| FALSE | 11 | 11 | 17 | 4 | 7 | 9 | 11 | 19 | 15 | 13 | 8 | 12 | 12 | 9 | 10 | 11 |
| Don't know | 23 | 24 | 12 | 41 | 28 | 21 | 22 | 16 | 15 | 22 | 26 | 26 | 24 | 20 | 25 | 22 |
| It is not uncommon for males to involuntarily ejaculate during a physical rectal exam of the prostate | | | | | | | | | | | | | | | | |
| TRUE | 18 | 18 | 4 | 17 | 22 | 21 | 17 | 16 | 16 | 19 | 18 | 22 | 14 | 19 | 20 | 22 |
| FALSE | 22 | 22 | 7 | 18 | 20 | 22 | 23 | 22 | 25 | 26 | 16 | 24 | 25 | 23 | 18 | 17 |
| Don't know | 60 | 59 | 89 | 64 | 58 | 57 | 60 | 62 | 59 | 55 | 66 | 54 | 61 | 58 | 62 | 61 |
| Prostate cancer is the most common cancer among males | | _ | | | | | | | | | | _ | | | | |
| TRUE | 54 | 54 | 47 | 45 | 49 | 56 | 52 | 58 | 61 | 56 | 51 | 53 | 54 | 56 | 53 | 51 |
| FALSE Don't know | 8 39 | 8 39 | 5 48 | 5 50 | 9 42 | 8 36 | 7 41 | 9 33 | 9 30 | 9 35 | 6 43 | 8 39 | 8 38 | 6 38 | 8 40 | 8 41 |

YouGov - Prostate cancer

Sample Size: 1456 adults in GB who were described as male at birth (i.e. who have a prostate)

| | | As far as you are prostate cane | | |
|---|-------|------------------------------------|---------------|------------------|
| | Total | A blood analysis test | A rectal exam | Both are typical |
| Weighted Sample | 1618 | 144 | 535 | 517 |
| Unweighted Sample | 1456 | 147 | 450 | 499 |
| | % | % | % | % |
| Vulnerability to prostate cancer is genetic i.e. you are more like to get prostate cancer if other members of your family got prostate cancer | | | | |
| TRUE | 35 | 36 | 34 | 47 |
| FALSE | 14 | 19 | 16 | 15 |
| Don't know | 52 | 45 | 51 | 38 |
| Prostate cancer requires immediate treatment once discovered | | - | | |
| TRUE | 66 | 65 | 72 | 71 |
| FALSE | 11 | 21 | 8 | 16 |
| Don't know | 23 | 14 | 20 | 13 |
| It is not uncommon for males to involuntarily | | | | |
| ejaculate during a physical rectal exam of the prostate | | _ | | |
| TRUE | 18 | 17 | 19 | 23 |
| FALSE | 22 | 26 | 26 | 26 |
| Don't know | 60 | 57 | 55 | 50 |
| Prostate cancer is the most common cancer among males | | _ | | |
| TRUE | 54 | 60 | 59 | 66 |
| FALSE | 8 | 8 | 7 | 9 |
| Don't know | 39 | 32 | 35 | 24 |